




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(15) International Patent Classification ⁶ : C07D 487/22		A1	(11) International Publication Number: WO 97/32885
		(43) International Publication Date: 12 September 1997 (12.09.97)	
(21) International Application Number: PCT/US97/03891		(81) Designated States: CA, JP, MX, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LC, MC, NL, PT, SE)	
(22) International Filing Date: 7 March 1997 (07.03.97)			
(30) Priority Data: 08/613,134 8 March 1996 (08.03.96) US Not furnished 6 March 1997 (06.03.97) US		Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(71) Applicant: HEALTH RESEARCH, INC. [US/US]; Roswell Park Memorial Institute Division, 666 Elm Street, Buffalo, NY 14263 (US).			
(72) Inventors: PANDEY, Ravindra, K.; 75 Lemay Court, Williamsville, NY 14221 (US). KOZYREV, Andrei, N.; 175 Cambridge Boulevard #1, Amherst, NY 14226 (US). DOUGHERTY, Thomas, J.; 2306 W. Oakfield, Grand Island, NY 14072 (US).			
(74) Agent: DUNN, Michael, L.; Dunn & Associates, P.O. Box 96, Newfane, NY 14108 (US).			

(54) Title: SYNTHESIS OF ISOIMIDE OF CHLORINS AND BACTERIOCHLORINS AND THEIR USE FOR DIAGNOSIS AND TREATMENT OF CANCER

(57) Abstract

Compounds having utility as light absorbing compounds, especially in the area of photodynamic therapy. Such compounds have formula (I), where $z = O$ or NR_{14} ; R_{14} is alkyl or substituted alkyl, R_1 is an amino acid group, a polyamine group, a polyether group or OR_{13} where R_{13} is alkyl; R_4 through R_{11} are $-H$, $-OH$, alkyl, alkylen, $-OR_{14}$ where R_{14} is H , alkyl or aryl, or a carbonyl containing group, provided that: R_4 may be taken together with R_5 to form $-O-$; R_6 may be taken together with R_7 to form $-O-$; R_8 may be taken together with R_9 to form $-O-$; R_{10} may be taken together with R_{11} to form $-O-$; and R_4 and R_7 may together form a chemical bond and R_8 and R_{11} may together form a chemical bond; and R_{12} is hydrogen or lower alkyl; provided that if one z is O , the other z is $-NR_{14}$.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	NL	Netherlands	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

SYNTHESIS OF ISOIMIDE OF CHLORINS AND BACTERIOCHLORINS AND
THEIR USE FOR DIAGNOSIS AND TREATMENT OF CANCER

Background of the Invention

This invention relates to treatment and diagnosis of cancer through the use of IR imaging and photodynamic therapy employing porphyrin related compounds and more particularly certain analogs of chlorins and bacteriochlorins.

Photosensitizers are chemicals which kill cells and/or fluoresce when activated by light of a specific wavelength. Most malignant and some premalignant tissues retain these photochemically active substances in higher concentrations and for longer durations than surrounding normal tissues. The retention time is not dependent on whether or not the cells are synthesizing DNA or undergoing cell growth or nutrient uptake.

Light sensitive drugs, lasers and fiber optic probes have been combined in a procedure known as photodynamic therapy (PDT). PDT has emerged as one of the most promising strategies in cancer treatment (including cancer detection). PDT is being increasingly used where chemotherapy, surgery and high energy irradiation have failed. In this new therapy, patients are given intravenous injections of a photodynamic drug that accumulates in cancer cells in much higher concentrations than in normal cells. The photodynamic (photosensitizing) drug is then activated to

kill the cancer cells by a laser beam directed to cancer cells through fiber optics. People who are too old or too frail to tolerate the stress of major surgery, chemotherapy or high energy radiation may be helped by photodynamic therapy, which often requires just local anesthesia and does not necessitate hospital admission. PDT is an important form of cancer therapy that has many additional advantages, e.g. it can be performed any number of times on a single patient, it is not contraindicated with other cancer therapies and it allows selective treatment of malignant tissues due to preferential retention of dye in cancer cells and it has already been established that for early or superficial malignancy, PDT may be curative.

Photosensitizers have been recognized for almost a century. In 1900, (Rabb, C., *Z. Biol.*, 1900, 39, 1423) reported the lethal effects of a combination of acridin orange dye and ordinary light on *Paramecium*. In 1903, von Tappeneir reported the first therapeutic use of photosensitizers when he used eosin and white light to treat skin tumors. The phototoxic effect of an administered porphyrin in man was observed in 1913. The localization of administered porphyrins in tumor tissue was recognized in the 1940's. It was not until 1972, however, that these two ideas (photodegradation of tissue and localization in tumors) came together successfully, when Diamond demonstrated that a porphyrin could preferentially degrade tumor implants in a rat (Diamond, I.; McDonagh, A.F.; Wilson, C.B.; Granelli, S.G.; Nielsen, S.; Jaenicke, R., *Lancet*, 1972, 1175). This result was confirmed and extended by Dougherty, T.J.; Grindey, G.B.; Fiel,

R.; Weishaupt, K.R.; Boyle, D.G.; *J. Natl. Cancer Inst.*, 1975, 55, 115.

The higher concentration of porphyrins in malignant tumors is used for the treatment and detection of cancer. For detection of early stage small tumors, the porphyrin-containing tumor cells and surrounding tissues are exposed to light. The porphyrins then emit a strong fluorescence, which contrasts with the much weaker fluorescence from the normal tissue, allowing for detection. For the treatment of cancer, photodynamic therapy (PDT) consists of injecting the patient with a photoactive dye and irradiating the tumor area with a wavelength of light which activates the dye to produce toxins which kill the tumor. The porphyrin dyes become toxic to the surrounding environment by producing single oxygen and oxygen radicals (Dougherty, T.J.; Kaufman, J.H.; Goldfarb, A.; Weishaupt, K.R.; Boyle, D.; Mittleman, A.; *Cancer Res.*, 1976, 38, 3628). PDT techniques depend strongly on how well the compound used preferentially concentrates within the tumor cell. Skin photosensitivity is the only known side effect of PDT with certain porphyrin type photosensitizers. Because skin retains these chemicals in enough quantities to produce surface reactions, patients must avoid exposure to sunlight.

The distribution of porphyrin drugs in the body compared with tumor cells is still under investigation. The distribution varies with cell type and porphyrin derivative. It is thought that once the photosensitizer is injected intravenously, some of

the drug escapes the blood stream and moves into the interstitial fluid. The porphyrin binds to the cellular membrane and slowly diffuses into the cell cytoplasm. Each porphyrin, then, rapidly binds to hydrophobic regions inside the cell. Fluorescence microscopy of porphyrin-treated leukemia L1210 cells shows a localization around the plasma membrane and within the intracellular vesicles.

Photofrin®, a hematoporphyrin derivative (Dougherty, T.J.; Boyle, D.G.; Weishaupt, K.R., "Photodynamic Therapy - Clinical and Drug Advances, Porphyrin Photosensitization," *Plenum Press*, New York, 1983, p. 3) is the only photosensitizer currently being used all over the world for the treatment of a variety of solid tumors. Hematoporphyrin derivative (Hpd) is prepared by mixing hematoporphyrin with glacial acetic acid and sulfuric acid, followed by hydrolysis and precipitation under acidic conditions. This method was partially described by Lipson et al (Lipson, R.L.; Baldes, E.J.; Olsen, A.M., *J. Natl. Cancer Inst.*, 1961, 26, 1). Hpd thus produced consists of a variety of porphyrins. When Hpd is separated into its two main fractions by gel filtration with Sephadex LH-20, the higher molecular weight portion, called Photofrin®, is a more efficient PDT agent (Dougherty, T.J.; Boyle, D.G.; Weishaupt, K.R.; Henderson, B.; Potter, W.; Bellnier, D.A.; Wityk, K.E., *Adv. Exp. Biol. Med.*, 1983, 160, 3). The recommended human dosage of Photofrin® is 1-2 mg/kg of body weight. The main components of Photofrin® are dimers and higher oligomers linked with ether, and possibly carbon-carbon linkages (Pandey, R.K.; Siegel, M.M.; Tsao, R.; McReynolds, J.M.;

Dougherty, T.J., *Biomed. and Environ. Mass Spectrometry*, 1990, 19, 405).

For a photosensitizer to be clinically useful, it must be non-toxic, selectively taken up and/or retained in malignant tissues, activated by penetrating light (>600 nm), and photochemically efficient. Although Photofrin® has been approved for commercialization in Canada, Europe and the United States, it lacks rapid clearance from tissues, is a complex mixture of oligomers, and has the disadvantage that its absorbance at 630 nm is not optimized for tissue penetration. New porphyrin photosensitizers are thus needed for the improvement of photodynamic therapy for cancer treatment.

There is a need for more efficient, chemically pure, less phototoxic, and better localizing porphyrins. Important prior art porphyrin and chlorin derivatives have been reviewed by Pandey, R.K.; Majchrzycki, D.F.; Smith, K.M.; Dougherty, T.J., *Proc. SPIE*, 1989, 1065, 104. The aspartyl derivatives of chlorin e_6 , monoaspartyl chlorin e_6 and diaspartyl chlorin e_6 , were found to be effective photosensitizers *in vitro* (Roberts, W.G.; Shaiu, F.Y.; Nelson, J.S.; Smith, K.M., Roberts, M.W., *J. Natl. Cancer Inst.*, 1988, 80, 330). With these compounds, the aspartyl group was noted to be responsible for the efficiency of tissue clearance. In pheophorbide, pyropheophorbide and chlorin e_6 series, certain alkyl ether derivatives including 2-(1-hexyloxyethyl)-2-des vinyl derivatives were found to be excellent photosensitizers compared with parent compounds, methyl

pheophorbide-a, pyropheophorbide- and chlorin₆. (Pandey, R.K.; Bellnier, D.A.; Smith, K.M.; Dougherty, T.J., *Photochem. Photobiol.*, 1991, 53, 65). This was attributed to the increased hydrophobicity of the hexyl group and is consistent with studies done by Evenson on porphyrins with varying polarities (Evenson, J.F.; Sommer, S.; Rimington, C.; Moan, J., *Br. J. Cancer*, 1987, 55, 483).

Chang, C.K.; Sotiroiu, C.; Wu, W., *J. Chem. Soc., Chem. Commun.*, 1986, 1213, have previously shown that chlorins, on reacting with osmium tetroxide can be converted to vic dihydroxy bacteriochlorin system. This methodology has been extended in the pheophorbide-a and chlorin *e*₆ series, by preparation at a series of vic -dihydroxy and keto-bacteriochlorins (Pandey, R.K.; Shiau, F.Y.; Sumlin, A.B.; Dougherty, T.J.; Smith, K.M., *Bioorg. & Med. Chem. Lett.*, 1992, 2, 491). It has also been reported that the regiospecificity of pyrrole subunits in osmium tetroxide oxidation is affected significantly by the presence of electron withdrawing substituents in the macrocycle(5a). These stable bacteriochlorins, prepared from mesochlorin *e*₆ trimethylester and pyropheophorbide-a methylester, have strong absorption in the red region (730 to 750 nm), but, did not show any significant *in vivo* photosensitizing activity (Kessel, D.; Smith, K.M.; Pandey, R.K.; Shiau, F.Y.; Henderson, B., *Photochem. Photobiol.*, 1993, 58, 200).

Hoober, J.K.; Sery, T.W.; Yamamoto, Y., *Photochem. Photobiol.*, 1988, 48, 579 showed that purpurin-18 2, which has

strong absorption at 700 nm might be a useful photosensitizer for photodynamic therapy (PDT).

Among long wavelength absorbing photosensitizers, bacteriochlorins have been proposed as potential useful candidates for use in photodynamic therapy (PDT) where strong absorptions in the visible spectrum can be used to photoactivate dyes previously located in targeted (neoplastic) tissues (Pandey, R.K.; Shiau, F.Y.; Isaac, M.; Ramaprasad, S.; Dougherty, T.J.; Smith, K.M., *Tetrahedron Lett.*, 1992, 33, 7815). Some naturally occurring bacteriochlorins, have previously been reported as effective photosensitizers both *in vitro* and as *in vivo* (Beems, E.M.; Dubbelman, T.M.A.R.; Lugtenburg, J.; Best, J.A.B.; Smeets, M.F.M.A.; Boehrgeim, J.P.J., *Photochem. Photobiol.*, 1987, 46, 639). However, most of the naturally occurring bacteriochlorins (760-780 nm) are extremely sensitive to oxygen, which results in rapid oxidation to the chlorin state (640 nm); thus the spectroscopic properties of the bacteriochlorins are lost. Further, if a laser is used to excite the bacteriochlorin *in vivo*, oxidation may result in the formation of a new chromophore absorbing outside the laser window, thus reducing the photodynamic efficiency.

It has been found that certain cyclic amide derivatives of porphyrins, including bacteriochlorins and chlorins, have both an increased wavelength and the requisite stability to meet the requirements of an improved photodynamic therapeutic agent. Unfortunately, the preparation of such compounds is difficult and yields are very low, e.g. 10 to 30%. Such compounds (e.g.

compound 8) have in the past been prepared by cyclizing chlorin-p₆ 6-N-hexylamide-7-methyl ester 3A (compound 3 reacted with diazomethane). The resulting cyclized reaction mixture contains a number of products in addition to the cyclic amine, especially unreacted starting material 3A. The synthesis route therefore is not only inefficient because of low yield, but requires significant subsequent purification. Further such compounds have not had optimal hydrophylic-lipophylic balance.

Brief Description of the Drawings

Figure 1 is a schematic equation showing the degeneration of bacteriochlorophyll-a to chlorin.

Figure 2 is a schematic equation showing the synthetic route to compounds 6 and 7 of the invention and their use as intermediates to compound 8.

Figure 3 is a schematic equation showing the synthetic route to cyclic anhydride, compound 1.

Figure 4 is a schematic equation showing the synthetic route to cyclic isoimides and cyclic imide.

Figure 5 is a schematic equation showing the synthetic route to isoimide bacteriochlorins 17 and 18 of the invention.

Figure 6A is a reflection spectroscopy curve showing preferential accumulation of compound 18 in tumors over muscle.

Figure 6B is a curve showing the ratio of tumor over muscle accumulation as represented in Figure 6A.

Brief Description of the Invention

In accordance with the present invention, there are provided new chlorin and bacteriochlorin derivatives having utility as fluorescent and photosensitizing compounds. Such compounds may be excited by microwaves, ultrasound, and visible or infrared radiation.


All of such novel compounds described herein may be used in traditional areas where compounds having such properties have utility. The compounds, may, for example, be incorporated into a substance such as a plastic product, excited with ultrasound, microwaves or visible light followed by using known methods for detecting emitted radiation to image the product for the purpose of detecting voids or other flaws in the product.

Certain of such compounds have special utility as photosensitizers in the area of photodynamic therapy for the detection and treatment of tumors.

In accordance with the invention, to make PDT more applicable, there is a need of long wavelength absorbing photosensitizers such as stable bacteriochlorins which have the ability to localize in high concentration at the tumor site.

Furthermore there is a need for an efficient and cost effective method for preparing such photosensitizers.

In accordance with the invention, a compound is therefore provided which comprises a chemical of the formula:



wherein z is $= O$ or $= NR_{14}$; R_1 is an amino acid group, a polyamine group, a polyether group or OR_{13} where R_{13} is alkyl; R_{14} is alkyl, or substituted alkyl, a polyamine group, or an amino acid group; R_4 through R_{11} are $-H$, $-OH$, alkyl, alkylene, $-OR_{16}$, where R_{16} is H , alkyl or aryl, or a carbonyl containing group, provided that; R_4 may be taken together with R_5 to form $=O$; R_6 may be taken together with R_7 to form $=O$; R_8 may be taken together with R_9 to form $=O$; R_{10} may be taken together with R_{11} to form $=O$; and R_4 and R_7 may together form a chemical bond and R_8 and R_{11} may together form a chemical bond; and R_{12} is hydrogen or lower alkyl; provided that if one z is $=O$, the other z is $=NR_{14}$.

The invention further includes a method for using the above compound as an intermediate for the preparation of additional

long wave length stable photosensitizers and a method for its preparation.

Detailed Description of the Invention

The invention permits more flexibility in the preparation of porphyrin-type compounds than was previously possible. Intermediate compounds may be provided with a number of substituents on the a and b rings and variable substituents at R_1 .

The a and b rings may be saturated or unsaturated at the R_4 - R_{11} positions or may contain hydrogen, hydroxy, formal or substituted and unsubstituted alkyl, alkoxy, alkenyl, aryl, and aryloxy groups. The alkyl, alkoxy, alkenyl, aryl and aryloxy groups usually contain 1 through 8 carbon atoms and more commonly contain 1 through 3 carbon atoms. A limited number, i.e., as many as 2, of such carbon containing groups may be long chain carbon containing groups, e.g., up to 22 carbon atoms.

The carbon containing groups may be substituted with carbonyl, hydroxy, phosphoro, carboxy, halo, sulfo, amino and ether substituents.

To obtain the compounds of the invention, a substituted or unsubstituted chlorin or bacteriochlorin is reacted by known methods, as described in Kenner et al., J. Chem. Soc., Perkin Trans. I, 1973, 2517, to obtain a six membered anhydride ring fused to the macrocycle. For example, to obtain a suitable bacteriochlorin, bacteriochlorophyll a is converted to bacteriopurpurin-a containing an anhydride ring before subjecting

it to the subsequent reactions described herein. The anhydride is then reacted with 1, hexyl amine to open the anhydride ring, e.g., as shown in Figure 2 to obtain 6 carboxylhexyl amide 2 and γ -carboxy-hexylamide 3.

The 6 carboxyhexylamide and the γ -carboxyhexylamide are then reacted separately or together with a carbodiimide to form a porphyrin diimide which is unstable and immediately converts to an isoimide, the compound of the invention. A preferred carbodiimide is dicycloberylcarbodiimide (DCC), which results in compounds similar to 6 and 7 in Figure 2 where the R_1 - R_{12} substituents may vary as described herein.

The invention may be described in more detail by reference to the following specific embodiment.

Initially, in order to establish the reaction conditions, purpurin-18 methyl ester 1 was used as the starting material. As expected, reaction of 1 (λ_{\max} 700 nm) with 1-hexylamine gave the corresponding amides in 95% yield as a mixture of 2 and 3 in the ratio of 9 to 1 (determined by using proton NMR) with λ_{\max} at 665 nm. Attempts to convert the amides 2 or 3 into the corresponding imides by following the methods used in converting aromatic anhydrides into imides, (e.g. such as heating with imidazole at 140°C) mainly gave decomposition products. Leaving the amide solution in dichloromethane or tetrahydrofuran at room temperature for a week gave a mixture of purpurins with cyclic anhydride 1 (700 nm), cyclic imide 8 (705 nm) in minor amounts and the starting material as a major product. By refluxing the

reaction mixture at various temperatures, slightly improved the yield of purpurin-anhydride without formation of any desired imide analogue. Various attempts were then made to optimize the reaction conditions, as summarized in Table 1. Reaction of amides 2 and 3 with K-10 clay using CH_2Cl_2 as a solvent again gave a mixture of cyclic imide 8 as minor product (12%) and anhydride analog 1 as major product (85%), which were easily separated by column chromatography. Reaction of purpurin amides (a mixture of 6-carboxyhexylamide 2 and γ -carboxy-hexylamide 3) with DCC afforded corresponding carbodiimides which are not stable and converted to corresponding isoimide analogues 6 and 7 (1:6) in 96% yield. This reaction is precisely the same reaction described in parent Application Serial No. 08/613,134, except that the useful end products were inadvertently mischaracterized as the unstable carbodiimide analogs rather than isoimides. Separation of the mixture gave pure isomers 6 and 7 with long wavelength absorptions at λ_{max} 696 and 690nm respectively. Both isomers were characterized by proton NMR and mass spectroscopy. Refluxing the isoimide (6 and 7) with various solvents alone or with K-10 clay gave mainly the starting material 1. Treatment of isoimides (either 6 or 7) with DBU/toluene at 60°C produced imide 8 in 60% yield. Interestingly, replacing DBU with stronger bases, such as methanolic KOH or NaOH at room temperature, gave the desired purpurin-imide in 85% overall yield (λ_{max} 705nm). This reaction was repeated several times using individual isoimide isomers (6 or 7), and produced the desired imide 4 in >80% yield without formation of purpurin-18 methyl ester 1.

Replacing DCC with 1,1'-thiocarbonyldiimidazole under similar reaction conditions gave purpurin-imide, but in a lower yield (Table 1).

A proposed mechanism for the formation of the imide ring in accordance with the invention is shown in Figure 2. In brief, addition of the carboxylic acid to the carbodiimide will give an O-acylisourea, an activated carboxylic acid derivative. It is now understood that these intermediate species are not stable and convert to corresponding isoimides and dicyclohexylurea. Intramolecular nucleophilic attack under basic conditions will generate cyclic imide. In tetrapyrrole chemistry, this is the first example of the formation of cyclic isoimides and imides from appropriate dicyclohexylcarbodiimide intermediate with methanolic KOH.

Related carbodiimide and isoimide analogs were prepared using bacteriopurpurin-a **12** as a substrate and converting it to the related imide derivative. Bacteriopurpurin-a **12** was isolated from *R. Spheroides* by following the methodology as described in U.S. Patent Application Serial No. 08/247,866 by reaction of bacteriochlorin with *n* propanol.

Another compound **12a** was obtained by treatment of bacteriochlorophyll-a from *R-speroides* with *n*-propanol. Reaction of **12a** with *n*-hexylamine gave the amide analogs **13** and **14**, which on reacting with dicyclohexylcarbodiimide is believed to produce the corresponding unstable carbodiimide derivatives **15** (minor component) and **16** (major component) which convert to

corresponding isoimides with long wavelength absorption at λ_{\max} 796nm 17 ($\epsilon=89,000$) and λ_{\max} 18 804 ($\epsilon=109,000$) respectively. Currently, the synthesis of a series of related compounds with various hydrophobic properties is in progress, e.g., (a) to increase/decrease the length of alkyl amides by opening the anhydride ring with various amines, and amino acids, (b) to replace the methylester group (at position -7, ring D) with various esters or aspartic acid analogs, (c) to replace dicyclohexylcarbodiimide with other carbodiimides, and (d) to introduce primary- or secondary alkyl ether groups at position 2- of the macrocycle.

For a compound to be useful for PDT and IR imaging, it should have preferential accumulation in tumor. In an initial experiment, the uptake of bacteriopurpurin 18 in tumor vs. muscle was measured by in vivo reflection spectroscopy. From Figures 6A and 6B it can be seen that bacteriopurpurin 18 shows preferential accumulation of drug in tumor than muscles (8:1). Studies with other related compounds are currently in progress.

Experimental

Chemistry:

Commercially available compounds and reagents were purchased from Aldrich, ACROS Organics and Sigma. Mps were taken using a Fisher-Johns hot plate melting point apparatus and are uncorrected. NMR spectra were recorded at 300 MHz on a GE instrument using $CDCl_3$ as solvent. Electronic absorption spectra were recorded using a Genesis-5 spectrophotometer.

New and novel method for the preparation of purpurin-imide from isoimide via carbodiimide intermediate:

Purpurin-18 methyl ester 1 (200 mg, 0.34 mmol) in dichloromethane (100 ml) was treated with 1-hexylamine (2 ml, 2 mmol) and stirred at room temperature for 20 h. Spectrophotometry was used to monitor disappearance of the peak at 700 nm and appearance of a new peak at 666 nm. The solvent was then removed under high vacuum, and the residue was crystallized from dichloromethane/hexane to give hexylamine derivatives 2 (major) and 3 (minor) as a mixture in 84 and 12% yields, respectively (total: 220 mg). The reaction mixture (220 mg, 0.34 mmol) was dissolved in dichloromethane (25 ml) and reacted with dicyclohexylcarbodiimide (DCC) (400 mg, 1.75 mmol) under a nitrogen atmosphere with stirring for 12 h. The solvent was concentrated to 10 ml and left overnight in the refrigerator; dicyclohexylurea formed as a by-product was removed by filtration. The filtrate was concentrated and separated into individual isomers 4 and 5 (in the ratio of 1 to 6) using preparative plates (silica gel). The structure of these compounds were determined to be isoimide derivatives. The intermediate carbodiimides 4 and 5 were not stable. The mechanism of the formation of these compounds is shown in Figs. 2 and 4. The yield was 90% (245 mg). For the preparation of purpurin-imide, the mixture of 6 and 7 (245 mg) was dissolved in THF (50 ml), and a methanolic solution of KOH (0.5 mg/10 ml water) was added. The reaction mixture was stirred for 5 min,

and was monitored by spectrophotometry (appearance of a new peak at 705 nm). The mixture was then diluted with dichloromethane (100 ml) and washed with water (3 x 100 ml). The organic layer was dried over anhydrous sodium sulfate. Evaporation of the solvent gave a residue which was chromatographed on a silica gel column (elution with 2% acetone/dichloromethane). The appropriate eluates were combined. The residue obtained after evaporating the solvent was crystallized from dichloromethane/hexane, and the desired purpurin-imide **8** was obtained in 85% yield (185 mg from **1**). For the syntheses of other purpurin-imides **9-11** (see Table 2), the intermediate amides, carbodiimides and isoimide analogs were not isolated, but were converted into the desired imides in one-pot without further purification.

Isoimide derivative of Bacteriopurpurin-a propyl ester (17) and (18):

Bacteriopurpurin-a propyl ester **12a** (30 mg, 0.05 mmol), obtained from *R. spheroides* was dissolved in dichloromethane (100 ml) and was treated with 1-hexylamine (0.2 ml, 0.1 mmol). The reaction mixture was stirred at room temperature for 24h; spectrophotometry was used to monitor the disappearance of the peak at 813nm (due to starting material) and appearance of a new peak at 786nm. The solvent was then removed under high vacuum, and the residue was crystallized from dichloromethane/hexane to give hexylamine derivatives **13** (minor) and **14** (major) as a mixture in 10 and 90% yields respectively. The reaction mixture

(30 mg, 0.05 mmol) was dissolved in dichloromethane (25 ml) and reacted with dicyclohexylcarbodiimide (DCC) (50 mg, 0.25 mmol) under a nitrogen atmosphere with stirring for 12 h. The solvent was concentrated to 10 ml and left overnight in the refrigerator; dicyclohexylurea formed as a by-product was removed by filtration. The filtrate was concentrated and separated into individual isomers 17 and 18 (in the ratio of 1 to 9) using preparative plates (silica gel). Yield: (28 mg, 72% overall). On the basis of NMR data, these compounds were found to be isoimide derivatives of bacteriopurpurin-a

Spectroscopic Data:

Purpurin-18-N-hexylimide Methyl Ester (8): [Fisher's Nomenclature]

Mp. 221-223°C. UV/Vis: (λ max/nm, ϵ): 705 (46,000); 647 (12,000); 549 (23,000); 510 (10,000); 483 (8,000); 417 (120,000). ^1H NMR (δ ppm, CDCl_3): 9.63 (s, β -meso H), 9.38 (s, α -meso H), 8.58 (s, δ -meso H), 7.92 (dd, J 19.5, 12.8 Hz, 2a-H), 6.29 (dd, J 19.5 Hz, 2b-H), 6.18 (dd, J 12.8 Hz, 2b'-H), 5.37 (d, J 8.5 Hz, 7-H), 4.48 (t, N-hexylimide-a- CH_2), 4.38 (q, J 8.0 Hz, 8-H), 3.84 (s, 5-Me), 3.62 (q, J 7.5 Hz, 4a- CH_2), 3.56 (s, OMe), 3.34 (s, 1-Me), 3.18 (3-Me), 2.65 (m, 7b-H), 2.51 (m, 7b'-H), 2.40 (m, 7a-H), 2.06 (m, 7a'-H), 2.00 (m, N-hexylimide-b,c- CH_2CH_2), 1.74 (d, J 8.0 Hz, 8-Me), 1.65 (t, J 7.2 Hz, 4-b Me), 1.43 (m, N-hexylimide-d,e- CH_2CH_2), 0.46 (t, J 7.8 Hz, N-hexylimide-f- CH_3), -0.08 and -0.17 (each br s, NH). m/z (LRMS): 661 (M+H).

Purpurin-18-15¹-hexylisoimide methyl ester (7)

Mp. 138-139°C. ¹H NMR (δ ppm, CDCl₃): 9.73 (s, β-meso H), 9.49 (s, α-meso H), 8.65 (s, δ-meso H), 7.92 (dd, J 19.4, 12.2 Hz, 2a-H), 6.31 (dd, J 19.4 Hz, 2b-H), 6.18 (dd, J 12.2 Hz, 2b'-H), 5.26 (d, J 8.5 Hz, 7-H), 4.57 (q, J 8.0 Hz, 8-H), 4.10 (m, hexyl isoimide-CH₂), 3.83 (s, 5-Me), 3.76 (q, J 7.5 Hz, 4a-CH₂), 3.58 (s, OMe), 3.46 (s, 1-Me), 3.28 (3-Me), 2.65 (m, 7b-H), 2.58-2.00 (m, 7b'-H; m, 7a-H; m, 7a'-H; m, hexyl isoimide-b,c-CH₂CH₂; 1.78 (d, J 8.0 Hz, 8-Me), 1.72 (t, J 7.2 Hz, 4-b Me), 1.61 (m, hexylamide-d,e-CH₂CH₂), 0.96 (t, J 7.8 Hz, hexyl isoimide f-CH₃), -0.61 and -0.88 (each br s, NH). m/z (LRMS): 661 (M + H).

Purpurin-18-13¹-hexyl isoimide methyl ester (6)

Mp. 142-143°C. ¹H NMR (δ ppm, CDCl₃): 9.74 (s, β-meso H), 9.57 (s, α-meso H), 8.75 (s, δ-meso H), 7.94 (dd, J 19.2, 12.5 Hz, 2a-H), 6.33 (dd, J 19.2 Hz, 2b-H), 6.15 (dd, J 12.5 Hz, 2b'-H), 5.24 (d, J 8.5 Hz, 7-H), 4.52 (q, J 8.0 Hz, 8-H), 4.06 (t, hexyl isoimide-a-CH₂), 3.81 (s, 5-Me), 3.74 (q, J 7.5 Hz, 4a-CH₂), 3.56 (s, OMe), 3.42 (s, 1-Me), 3.24 (3-Me), 2.65 (m, 7b-H), 2.51-2.00 (m, 7b'-H; m, 7a-H; m, 7a'-H; m, hexylisoimide-b,c-CH₂CH₂; m, dicyclohexylisourea-CH₂), 1.76 (d, J 8.0 Hz, 8-Me), 1.68 (t, J 7.2 Hz, 4-b Me), 1.58 (m, hexyl isoimide-d,e-CH₂CH₂), 0.98 (t, J 7.8 Hz, hexylisoimide-f-CH₃), -0.66 and -0.84 (each br s, NH). m/z (LRMS): 661 (M+H).

Purpurin-18-N-hexylimide-7-N-aspartylamide Dimethyl Ester (9):
[Fisher's Nomenclature]

Mp. 218-219°C. UV/Vis: (λ max/nm, ϵ): 705 (43,200); 648 (11,000); 549 (21,000); 510 (9,200); 483 (7,800); 417 (112,000). ^1H NMR (δ ppm, CDCl_3): 9.60 (s, β -meso H), 9.34 (s, α -meso H), 8.58 (s, δ -meso H), 7.90 (dd, J 19.8, 12.6 Hz, 2a-H), 6.98 (d, J 9.6, aspartate-NHCO), 6.32 (dd, J 19.8 Hz, 2b-H), 6.18 (dd, J 12.6 Hz, 2b'-H), 5.34 (d, J 8.5 Hz, 7-H), 4.46 (t, N-hexylisoimide-a- CH_2), 4.44 (q, J 8.0 Hz, 8-H), 4.38 (m, aspartate-CH), 3.82 (s, 5-Me), 3.69 (s, OMe), 3.64 (q, J 7.5 Hz, 4a- CH_2), 3.61 (s, OMe), 3.36 (s, 1-Me), 3.16 (3-Me), 2.84 (m, aspartate- CH_2), 2.64 (m, 7b-H), 2.51 (m, 7b'-H), 2.46 (m, 7a-H), 2.06 (m, 7a'-H), 1.99 (m, N-hexylimide-b,c- CH_2CH_2), 1.75 (d, J 8.0 Hz, 8-Me), 1.66 (t, J 7.5 Hz, 4-b Me), 1.45 (m, N-hexylimide-d,e- CH_2CH_2), 0.96 (t, J 7.8 Hz, N-hexylimide-f- CH_3), -0.38 and -0.11 (each br s, NH). m/z (LRMS): 791 (M+H).

Purpurin-18-N-hexylimide-7-N-Aspartylamide-di-tert-butyl Ester
(10):

Mp. 190-192°C. UV/Vis: (λ max/nm, ϵ): 705 (42,800); 648 (11,000); 549 (20,000); 510 (9,000); 483 (7,500); 417 (110,000). ^1H NMR (δ ppm, CDCl_3): 9.63 (s, β -meso H), 9.37 (s, α -meso H), 8.58 (s, δ -meso H), 7.86 (dd, J 19.4, 12.5 Hz, 2a-H), 6.74 (d, J 9.6 Hz, aspartate-NHCO), 6.32 (dd, J 19.4 Hz, 2b-H), 6.21 (dd, J 12.5 Hz, 2b'-H), 5.33 (d, J 8.5 Hz, 7-H), 4.66 (q, J 8.0 Hz, 8-H), 4.41 (t, N-hexylimide- CH_2), 3.99 (m, aspartate-CH), 3.84 (s, 5-Me), 3.68 (q, J 7.5 Hz, 4a- CH_2), 3.36 (s, 1-Me), 3.18 (3-Me),

2.76 (m, aspartate-CH₂), 2.65 (m, 7b-H), 2.51 (m, 7b'-H), 2.46 (m, 7a-H), 2.06 (m, 7a'-H), 1.92 (m, N-hexylimide-b,c-CH₂CH₂), 1.68 (d, J 8.0 Hz, 8-Me), 1.66 (t, J 7.5 Hz, 4-b Me), 1.37 (s, aspartate-t Bu), 1.34 (s, aspartate-t Bu), 1.14 (m, N-hexylimide-d,e-CH₂CH₂), 0.95 (t, J 7.8 Hz, N-hexylimide-f-CH₃), -0.38 and -0.11 (each br s, NH). Mass: m/z (HRMS): requires for C₅₁H₆₇N₆O₇: 875.5071. Found 875.5016.

Purpurin-18-N-glycylimide-tert-Butyl Ester-7-aspartylamide-di-tert-butyl Ester (11): [Fisher's Nomenclature]

Mp. 138-139°C. UV/Vis: (λ max/nm, ε): 705 (41,300); 649 (10,000); 549 (19,000); 510 (8,700); 483 (7,300); 417 (105,000). ¹H NMR (δ ppm, CDCl₃): 9.61 (s, β-meso H), 9.35 (s, α-meso H), 8.56 (s, δ-meso H), 7.88 (dd, J 19.5, 12.6 Hz, 2a-H), 6.65 (d, J 9.5 Hz, aspartate-NHCO), 6.26 (dd, J 19.5 Hz, 2b-H), 6.16 (dd, J 12.5 Hz, 2b'-H), 5.27 (d, J 8.5 Hz, 7-H), 5.18 (q, glycine-CH₂), 4.68 (q, J 7.5 Hz, 8-H), 4.38 (m, aspartate-CH), 3.38 (s, 5-Me), 3.64 (q, J 7.5 Hz, 4a-CH₂), 3.35 (s, 1-Me), 3.16 (3-Me), 2.79 (m, aspartate-CH₂), 2.66 (m, 7b-H), 2.54 (m, 7b'-H), 2.46 (m, 7a-H), 2.01 (m, 7a'-H), 1.73 (d, J 7.5 Hz, 8-Me), 1.67 (t, J 7.5 Hz, 4-b Me), 1.58 (s, glycine-t Bu), 1.38 (s, aspartate-t Bu), 1.34 (s, aspartate-t Bu), 0.10 and -0.04 (each br s, NH). m/z (LRMS) 905.4 (M+H).

Bacteriopurpurin-a 17-propyl Ester (12a): [Fisher's Nomenclature]

UV/Vis (CH₂Cl₂, λmax, nm): 813 (56,000); 543 (32,000); 408 (48,000); 363 (102,000). NMR (CDCl₃, δ ppm): 9.21 (s, 1H, 5-H), 8.79 (s, 1H, 10-H), 8.62 (s, 1H, 20-H), 5.14 (d, 1H, J=8.0, 17-

H), 4.30 (m, 2H, 3-H, 18-H), 4.08 (m, 1H, 8-H), 3.94 (t, 2H, CO₂CH₂), 3.66 (s, 3H, 12-Me), 3.55 (s, 3H, 2-Me), 3.17 (s, 3H, 3-Me), 2.73 (m, H, 17b-H), 2.41 (m, 5H, CH₂CH₂CH₃ + 8a-CH₂ + 7b'-H), 2.14 (m, H, 17a-H), 1.98 (m, H, 17a'-H), 1.81, 1.73 (each d, 3H, J = 8.0, 18-Me, 7-Me), 1.11 (t, 3H, J = 7.8, 3-b Me), 0.83 (t, 3H, J = 8.2, CH₂CH₂CH₃), -0.30 and -0.67 (each br s, 2H, 21, 23-NH).

Bacteriopurpurin-a-15¹-N-hexylisoimide (17): [Fisher's Nomenclature]:

UV/Vis: λ_{max}: 795 (67,000); 537 (24,500); 410 (50,400); 363 (89,600). NMR (CDCl₃, δ ppm): 9.21 (s, 1H, 5-H), 8.77 (s, 1H, 10-H), 8.68 (s, 1H, 20-H), 5.35 (m, 2H, NHCO + 17-H), 4.32 (m, 2H, 3-H, 18-H), 4.11 (m, 3H, 8-H + hexylamide-a-CH₂), 3.91 (t, 2H, CO₂CH₂), 3.69 (s, 3H, 12-Me), 3.58 (s, 3H, 2-Me), 3.19 (s, 3H, 3-Me), 2.62 (m, H, 17b-H), 2.44 (m, 5H, CH₂CH₂CH₃ + 8a-CH₂ + 7b'-H), 2.14 (m, 6H, 17a-H + hexylisoimide-b,c-CH₂ + 17a'-H), 1.93, 1.84 (each d, 3H, J = 8.0, 18-Me, 7-Me), 1.60 (m, 4H, hexylamide-d,e-CH₂), 1.11 (t, 3H, J = 7.8, 3-b Me), 0.97 (t, 3H, hexylamide-f-CH₃), 0.88 (t, 3H, J = 8.2, CH₂CH₂CH₃), -0.68 and -1.03 (each br s, 2H, 21, 23-NH). Mass: LRMS:708(M+H)

Bacteriopurpurin-a-13¹-N-hexyl isoimide (18) [Fisher's Nomenclature]

UV/Vis (λ max/nm, ε): 804 (82,800); 539 (33,200); 409 (59,400); 360 (94,000). NMR (CDCl₃, δ ppm): d H: 9.38 (s, 1H, 5-H), 8.88 (s, 1H, 10-H), 8.73 (s, 1H, 20-H), 5.46 (m, 1H, NHCO), 5.18 (d,

1H, $J = 8.0$, 17-H), 4.34 (m, 2H, 3-H, 18-H), 4.17 (m, 1H, 8-H), 3.91 (m, 2H, hexylisoimide-a-CH₂), 4.06 (t, 2H, CO₂CH₂), 3.68 (s, 3H, 12-Me), 3.59 (s, 3H, 2-Me), 3.19 (s, 3H, 3-Me), 2.73 (m, H, 17b-H), 2.42 (m, 5H, CH₂CH₂CH₃ + 8a-CH₂ + 7b'-H), 2.14 (m, H, 17a-H), 2.08 (m, 5H, hexylamide-b,c-CH₂+17a'-H), 2.01, 1.93 (each d, 3H, $J=8.0$, 18-Me, 7-Me), 1.57 (m, 4H, hexylisoimide-d,e-CH₂), 1.12 (t, 3H, $J = 7.8$, 3-b Me), 0.96 (t, 3H, hexylamide-f-CH₃), 0.87 (t, 3H, $J = 8.2$, CH₂CH₂CH₃), -0.86 and -1.13 (each br s, 2H, 21, 23-NH). Mass: LRMS:708(M+1)

Biological Studies:

Determination of drug uptake:

The absorption spectrum of a compound in living tissue can be recorded using an instrument and technique which we have developed. The experiment measures the light which scatters through the tissue. The light originates in a high pressure Xenon arc lamp and passes through a grating monochromator to a 90 Hz chopper and then into a 400 μ m diameter quartz fiber. The distal end of this fiber is placed in contact with the tissue (e.g., an experimental mouse tumor) and the light is collected by a second fiber placed in contact with the tissue at a fixed distance (3 to 5 mm) from the first fiber. The noninvasive character of this measurement makes data collection possible at a number of time points after the i.v. injection of an experimental light absorbing compound (e.g., a potential photosensitizer). The light signal is detected by a photodiode. The photo current is converted into a voltage, amplified by a tuned (90 Hz)

amplifier and synchronously detected. The chopping at 90 Hz and synchronous detection makes examining spectra under normal room lighting possible.

For these experiments, the mice were first anesthetized using either Pentobarbital or Ketamine Xylazine i.p. The optical power as a function of wavelength was recorded before the i.v. injection of the sensitizer. The monochromator is set to the expected longest wavelength of the experimental drug's *in vitro* absorption spectrum. The drug was then administered by tail vein injection and the light signal recorded. The second spectrum contains a component caused by the presence of the drug in the tumor. This *in vivo* drug absorption spectrum is best displayed by taking the ratio of the post-injection spectrum to the pre-injection spectrum. This ratio offers the same advantages as a double beam absorption spectrophotometer. The pre-injection mouse data can be thought of as the reference beam sample (typically a cuvette and solvents) and the post-injection data as the sample beam containing everything in the reference beam plus the experimental drug. The ratio of these two spectra is certainly not influenced by the wavelength dependence of the light signal which characterizes the instrument. As a safeguard against day to day or hour to hour drift in the total light output of the lamp, both spectra (pre- and post-injection) are normalized by dividing the signal strength at a wavelength where the drug absorption is negligible.

Photosensitizing Efficacy: Part I - In vivo Efficacy of Bacteriochlorins 17 and 18 using SMT-F Tumor Model

The new photosensitizer was screened in a mouse/tumor model system. A model system consisted of observing the size reduction of the SMT-F tumor, a fast growing spontaneous mouse mammary tumor subline, transplanted subcutaneously to male or female DBA/2 HA-DD mice. The tumor line was maintained in vivo via serial transplantation in the same mouse strain. DBA/2 Ha-DD mice are readily available and were obtained locally.

When mice were both the appropriate age (approximately 6 weeks) and weight (approximately 20 g), small pieces of tumor (1-2 mm cube) were transplanted with a 18 gauge trocar from a donor tumor to recipient mouse. This technique provides for relatively uniform tumor size and allows location of the tumor in the right auxiliary region of the animal within each experimental group. Only animals with single tumors were chosen for experiments. When tumor reached 4-5 mm in diameter, the animals were injected with the potential photosensitizer chosen from the group described above. Prior to irradiation, the fur overgrown and surrounding the tumor was removed with electric clippers. Three or twenty four hours after injecting the drug, the mouse was placed in a custom-made aluminum holder.

Irradiation Conditions:

Standard light dose was 75 mW/cm^2 for 30 min for a total incident dose of 135 J/cm^2 from a tunable dye laser tuned to the maximum red absorption peak. Spectra Physics 2040, a quartz

fiber fitted with a microlens was interfaced to the dye laser deliver a uniform field of light. Laser output was measured with a power meter. Further studies at various light doses and treatment conditions are currently in progress.

Experimental Procedure:

Following light exposure, the mice were kept in groups of 5 per cage and supplied with pelleted food and tap water ad libitum. Tumor size and gross appearance of both tumor and overlying surrounding skin was monitored daily for 80 days after photoillumination unless growth of non-responsive tumor require early sacrifice of those animals.

The photosensitizer was dissolved in known quantity of Tween 80 (Aldrich) surfactant and diluted by a factor of 10 with saline solution to produce a final Tween 80 concentration of 10-20%. The solution was then filtered through a syringe filter. The concentration of the solution was determined on the basis of the extinction coefficient value of the photosensitizer at the longest wavelength absorption. Absorption spectra were obtained using a Spectronic Genesis5 spectrophotometer.

Before injecting the drug into mice, the purity of the compounds was ascertained by analytical HPLC using Spectra Physics HPLC, connected with C8 reverse phase column, eluted with methanol/water by adjusting the pH to 7.0 using phosphate buffer.

Alkylimide derivatives of bacteriochlorins:

Procedures similar to preparation of N-hexylimide derivatives were followed except that various other alkyl amines were substituted for hexylamine.

The resulting compounds were tested for *in vivo* antitumor activity.

4-6 mm diameter tumors (DBA/2 mice transplanted with SMT-F tumors) were exposed to 75 mW/cm² for 30 minutes to deliver light (135 J/cm²) from a tunable dye laser tuned to a maximum red absorption peak. The tumors were non-palpable and five mice were used per group.

The results are in Table 4. The results indicate a large suppressive effect for isomers where the extended alkyl group is closest to the amino acid group.

Part II - *In vivo* efficacy of bacteriochlorins 17 and 18 using Radiation Induced Fibrosarcoma (RIF) Tumor Model.

A mixture of bacteriochlorins 17 and 18 was also evaluated for *in vivo* PDT efficacy using another model (RIF tumor model) routinely used in our laboratory. In brief, six mice per group with appropriate tumor size (4-5 mm) were chosen for the experiment. As shown in Table 3, the animals were treated at variable doses of light and drug. Beginning 24h after PDT, and at least every other day thereafter, tumors were measured in orthogonal diameters with an electronic caliper (ultra-Cal Mark III; Fred V. Fowler Co., Boston MA). Each measurement was automatically recorded, where the tumor volume, V, was calculated


using the formula $v=(lw^2)/2$, where l is the longest axis of the tumor and w is the axis perpendicular to l . The time for growth to the tumor to 400 mm^3 was then estimated by interpolation of the times just before and after 400 mm^3 was reached. Tumor response was recorded on the basis of the number of animals which were found to be tumor free. Appropriate controls were carried out with tumor-bearing mice which received no treatment at all, or received light or photosensitizer only.

Table 1. Formation of purpurin-18 with cyclic anhydride and imide rings at various reaction conditions

Method No	Conditions	Yield (%)		
		imide 4	anhydride 1	starting materials (5 & 6)
1.	THF, refluxing for 4h.	5	70	25
2.	Imidazole, 140°C, 1h	Decomposition products		
3.	K-10 Clay, CH ₂ Cl ₂ , 24h	10-12	80-85	0
4.	CH ₂ Cl ₂ , 10 days	15-20	20-25	50
5. DCC with:				
	a. K-10 Clay, RT	0	0	100
	b. DBU, RT	0	0	100
	c. DBU, toluene, reflux 2h	60	0	40
	d. KOH/MeOH			
	RT, 10 min	85	0	0
6. TCD with:				
	a. DBU, RT	0	0	100
	b. DBU, toluene, reflux 2h	10	50	-
	c. DBU, THF 60°C	30	0	60
	d. KOH/MeOH			
	RT, 10 min	10	0	90

DCC: Dicyclohexylcarbodiimide; DBU, 1,8-Diazabicyclo(5.4.0)undec-7-ene; RT, Room temperature; TCD, 1,1'-Thiocarbonyldiimidazole.

Table 2. Yields of purpurin-imides




Compound No.	R	R ₁	% yield
8	OMe	n-hexyl	85
9	Asp-di-methyl ester	n-hexyl	89
10	Asp-di-tert-butyl ester	n-hexyl	87
11	Asp-di-tert-butyl ester	tert-butyl Gly	78

Table 3. PDT Efficacy in Mice Transplanted with Radiation Induced Fibrosarcoma (RIF) Tumor with a Mixture of Bacteriochlorins 17 and 18


Drug Dose $\mu\text{mol/kg}$	Light Dose Rate mW/cm^2	Timer after Injection hrs	Tumor Response days		
			1-2	7	14
0.20	75	24	100%	100%	80%
0.20	30	24	80%	60%	40%
0.25	75	24	100%	100%	80%
0.25	30	24	80%	80%	60%

TABLE 4



MINOR (795 NM)

17. R = (CH₂)₅CH₃
 19. R = (CH₂)₆CH₃
 21. R = (CH₂)₂CH₃
 23. R = (CH₂)₉CH₃



(MAJOR (804 NM)

18. R = (CH₂)₅CH₃
 20. R = (CH₂)₆CH₃
 22. R = (CH₂)₂CH₃
 24. R = (CH₂)₉CH₃

Comparative *in vivo* Antitumor Activity of Bacteriochlorins

Compound	Dose (μmol/kg)	in vivo absorption (λ max)	Time(h) betw. injection and light treatment	Tumor response (d) ^{†,*}		
				1-2	15	30 +
Mixture of 17 and 18	0.47	804	24	REGROWTH ON DAY 15		
20	0.47	804	24	NO RESPONSE		
22	0.47	804	24	REGROWTH ON DAY 4		
Mixture of 23 and 24	0.47	804	24	100	40	20

[†] 4-5 mm diameter tumors were exposed to 75 MW/cm² for 30 min to deliver 135 J/cm² light from a tunable dye laser tuned to the maximum red absorption peak.

d= days.

* Non-palpable tumors.


WHAT IS CLAIMED IS:

1. A method for the manufacture of an imide derivative of purpurin comprising:

reacting hexylamine with a chlorin or bacteriochlorin having a macrocycle with a six membered anhydride ring fused thereto, said macrocycle containing a and b rings which may be saturated or unsaturated at R₄ to R₁₁ positions of the rings and which R₄ and R₁₁ positions may contain at least one group selected from the group consisting of hydrogen, hydroxy, formal, substituted and unsubstituted alkyl, alkoxy, alkenyl, aryl and aryloxy wherein carbon containing groups may be substituted with a substituent selected from carboxyl, hydroxy, phosphoro, carboxy, halo, sulfo, amino and ether, to obtain a purpurin derivative; and reacting the purpurin derivative with a carbodiimide to obtain the imide derivative of purpurin.

2. The method of Claim 1, wherein the carbodiimide is dicycloberylcarbodiimide.

3. The method of Claim 1, wherein the imide derivative is further reacted with an alkali metal hydroxide to obtain a purpurin imide of the formula:



33

where R is selected from the group consisting of OMe and Asp-dimethyl ester and R_1 is selected from the group consisting of n-hexyl and tert-butyl Gly.

4. The method of Claim 1, wherein the imide derivative is further reacted with an alkali metal hydroxide to obtain a cyclic imide.


5. An imide of purpurin manufactured in accordance with the method of Claim 1.

6. An imide of purpurin manufactured in accordance with the method of Claim 2.

7. A reaction product comprising an imide of purpurin manufactured in accordance with the method of Claim 3.

8. A reaction product comprising an imide of purpurin manufactured in accordance with the method of Claim 4.


9. A compound of the formula:



34


where z is = 0 or = NR_{14} ; R_1 is an amino acid group, a polyamine group, a polyether group or OR_{13} where R_{13} is alkyl; R_{14} is alkyl, substituted alkyl, a polyamine group, or an amino acid group; R_4 through R_{11} are -H, -OH, alkyl, alkylene, - OR_{16} , where R_{16} is H, alkyl or aryl, or a carbonyl containing group, provided that; R_4 may be taken together with R_5 to form =O; R_6 may be taken together with R_7 to form =O; R_8 may be taken together with R_9 to form =O; R_{10} may be taken together with R_{11} to form =O; and R_4 and R_7 may together form a chemical bond and R_8 and R_{11} may together form a chemical bond; and R_{12} is hydrogen or lower alkyl; provided that if one z is = 0, the other z is = NR_{14} .

10. The compound of Claim 9 wherein R_{11} and R_{12} are $-\text{CH}_3$.
11. The compound of Claim 9 wherein R_2 is an allyl group.
12. The compound of Claim 9 wherein R_3 is an allyl group.
13. The compound of Claim 9 having the formula:




where R is normal alkyl of 2 through 12 carbon atoms.

14. The compound of Claim 9 having the formula:



where R is normal alkyl of 2 through 12 carbon atoms.

15. The compound of Claim 10 wherein R_2 is $O=COR_{15}$.
16. The compound of Claim 10 wherein R_3 is $O=COR_{15}$.
17. The compound of Claim 15 wherein R_1 is H_3CO_2C- .
18. The compound of Claim 16 wherein R_2 is H_3CO_2C- .
19. A method for the manufacture of an isoimide derivative of purpurin comprising reacting a purpurin of the formula:



wherein R_1 is an amino acid group, a polyamine group, a polyether group or OR_{13} where R_{13} is alkyl; R_{14} is alkyl, or substituted alkyl, a polyamine group, or an amino acid group; R_4 through R_{11} are -H, -OH, alkyl, alkylene, $-OR_{16}$, where R_{16} is H, alkyl or aryl, or a carbonyl containing group, provided that; R_4 may be taken together with R_5 to form =O; R_6 may be taken together with R_7 to form =O; R_8 may be taken together with R_9 to form =O; R_{10} may be taken together with R_{11} to form =O; and R_4 and R_7 may together form a chemical bond and R_8 and R_{11} may together form a chemical bond; and R_{12} is hydrogen or lower alkyl; provided that if one z is =O, the other z is $=NR_{14}$, with 1-hexylamine to open the anhydride ring followed by reaction with a carbodiimide to obtain the isoimide derivative of purpurin.

20. A method for the manufacture of purpurin imide comprising reacting the compound of Claim 9 with alkali metal hydroxide.

21. A method for the manufacture of purpurin imide comprising reacting the compound of Claim 10 with an alkali metal hydroxide.

22. A method for the manufacture of purpurin imide comprising reacting the compound of Claim 15 with an alkali metal hydroxide.

23. A method for the manufacture of purpurin imide comprising reacting the compound of Claim 16 with an alkali metal hydroxide.

24. A method for the manufacture of purpurin imide comprising reacting the compound of Claim 17 with an alkali metal hydroxide.

25. A method for the manufacture of purpurin imide comprising reacting the compound of Claim 18 with an alkali metal hydroxide.

26. A method for the manufacture of purpurin imide comprising reacting the purpurin derivative from Claim 19 with alkali metal hydroxide.

27. A method for the manufacture of purpurin imide comprising reacting the purpurin derivative from Claim 19 with alkali metal hydroxide.

28. A method for the manufacture of an imide derivative of purpurin comprising:

reacting hexylamine with a chlorin or bacteriochlorin having a macrocycle with a six membered anhydrid ring fused thereto, said macrocycle containing a and b rings which may be saturated or unsaturated at R_4 to R_{11} positions and which R_4 to R_{11} positions may contain at least one group selected from the group consisting of hydrogen, hydroxy, formal, substituted and unsubstituted alkyl, alkoxy, alkenyl, aryl and aryloxy, wherein carbon containing groups may be substituted with a substituent selected from carbonyl, hydroxy, phosphoro, carboxy, halo, sulfo, amino and ether, to obtain a purpurin derivative; and

reacting the obtained purpurin derivative with a carbodiimide to obtain the imide derivative of purpurin.

CONVERSION OF BACTERIOCHLOROPHYLL-A INTO CHLORIN
IN PRESENCE OF LIGHT AND AIR




FIG. 1





FIG. 2

3/7



FORMATION OF ANHYDRIDE RING

FIG 3




FIG. 4




FIG. 5

6/7




FIG 6A




FIG 6B

INTERNATIONAL SEARCH REPORT

International application No
PCT/US97/03891

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C07D 487/22

US CL : 540/145,472,474

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 540/145,472,474

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X, P	US 5,591,847 A (PANDEY et al) 07 JANUARY 1997 see entire document.	1-28
X	LEE et al. Use of the Chlorophyll Derivate, Purpurin-18, for Syntheses of Sensitizers for Use in Photodynamic Therapy., Journ. Chem. Soc. Perkin Trans I. October 1993, pages 2369-2372, especially page 2369 and formulae on page 2370.	1-28
Y	FIESER AND FIESER, Reagents for Organic Synthesis, Vol I, 1967, page 233. N.Y. John Wiley & Sons.	1-4 and 20-28.

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents	T later document published after the international filing date or priority date and not in conflict with the application but cited to underlain the principle or theory underlying the invention
A* document defining the general state of the art which is not considered to be of particular relevance	X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E* earlier document published on or after the international filing date	Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reasons (as specified)	
O* document referring to an oral disclosure, use, exhibition or other means	
P* document published prior to the international filing date but later than the priority date claimed	G* document member of the same patent family

Date of the actual completion of the international search

02 JUNE 1997

Date of mailing of the international search report

24 JUL 1997

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Authorized officer

PAVANARAM SRIPADA

Facsimile No (703) 305-3230

Telephone No (703) 308 4717

Form PCT/ISA/210 (second sheet) (July 1992)*